


**Juneberry or
Saskatoon
Serviceberry**



Juneberry or Saskatoon Serviceberry (*Amelanchier alnifolia*)

General Description

A hardy, medium to tall, suckering shrub, native to prairie hillsides and woody draws. Also called Saskatoon and Serviceberry. Fruits are highly prized for food. Selected cultivars are used in commercial production.

Leaves and Buds

Bud Arrangement - Alternate.

Bud Color - Chestnut-brown to purplish.

Bud Size - Medium, 1/4 to 3/8 inch.

Leaf Type and Shape - Simple, broad-oval.

Leaf Margins - Sometimes serrate-dentate, mostly above the middle, with rounded tips and bases.

Leaf Surface - Pubescent, becoming smooth.

Leaf Length - 1 to 2 inches.

Leaf Width - 3/4 to 1½ inches.

Leaf Color - Young leaves at first grayish-pubescent, soon becoming smooth and dark green; yellow fall color.

Flowers/Fruits

Flower Type - Erect racemes, at the tips of branches.

Flower Color - White.

Fruit Type - Berry-like pomes, with 4 to 10 small seeds.

Fruit Color - Purplish to black, with a leathery coat.

Form

Growth Habit - Upright.

Texture - Medium-fine, summer; medium-fine, winter.

Crown Height - 6 to 15 feet.

Crown Width - 5 to 12 feet.

Bark Color - Light brown to gray on older stems.

Root System - Shallow.

Environmental Requirements

Soils

Soil Texture - Prefers loam to sandy loam soil high in organic matter.

Soil pH - 5.5 to 7.5.

Windbreak Suitability Group - 1, 3, 4, 4C.

Cold Hardiness

USDA Zone 2.

Water

Needs adequate moisture to bear fruit. Limited drought tolerance, does not withstand ponding.

Light

Full sun to partial shade.

Uses

Conservation/Windbreaks

Large shrub for farmstead windbreaks, and riparian plantings.

Wildlife

High quality plant for wildlife cover and food. Stems, twigs, and leaves are browsed by deer and fruits are eaten by a variety of birds.

Agroforestry Products

Wood - Used as arrow shafts by Native Americans.

Food - Nutritious fruit used fresh, frozen or processed.

Medicinal - Crude extract used in cancer research.

Urban/Recreational

Good for naturalizing, screen plantings and landscape borders.

Cultivated Varieties

Superior quality, larger fruiting cultivars include:

Honeywood Juneberry (*Amelanchier alnifolia* 'Honeywood')

Northline Juneberry (*A. alnifolia* 'Northline')

Pembina Juneberry (*A. alnifolia* 'Pembina')

Smokey Juneberry (*A. alnifolia* 'Smokey')

Success Juneberry (*A. alnifolia* 'Success')

Related Species

Alleghany Serviceberry (*Amelanchier laevis*) - Taller growing species with potential windbreak and landscape value for the Northern Plains.

Pests

Intermittent flower or fruit loss may be caused by insects or late frosts.

SASKATOON

Amelanchier alnifolia (Nutt.)
Nutt. ex Roemer
Plant Symbol = AMAL2

Contributed By: USDA NRCS National Plant Data Center & the Biota of North America Program



William R. Hewlitt
© California Academy of Sciences
@ CalPhotos

Alternate common names

Saskatoon service-berry, service-berry, juneberry, shadbush

Uses

Saskatoon is planted as an ornamental and to produce commercial fruit crops. Many cultivars are commercially available, selected for desirable plant and/or fruit characteristics. Much research and literature details the development of cultivars and cultivation techniques.

The fruits are used in pies, jams, and fruit rolls and for making jelly and syrup. Saskatoon wine is a regional specialty. Native Americans ate the berries fresh and dried, often mixed with other foods for sweetening and flavor. Dried and rehydrated berries were added to dried vegetables and cooked into soups and puddings.

Native Americans boiled branches to make a tea for treating colds. A drink was made from the bark for

stomach problems. Bark and twigs provided a medicine for recovery after childbirth. In

combination with other plants, it was used to make a contraceptive. The strong and straight-grained wood was used to make arrows, digging sticks, spear shafts, tool handles, and seed beaters. Young branches were twisted into a type of rope.

Saskatoon is attractive as an ornamental shrub or may be trimmed as a hedge. It is an important species for reclamation, wildlife, watershed, and shelterbelt plantings. It can be started from seed or vegetative cuttings.

Status

Please consult the PLANTS Web site and your State Department of Natural Resources for this plant's current status, such as, state noxious status and wetland indicator values.

Description

General: Rose Family (Rosaceae). Native shrubs or small trees growing to 7 meters high, variable in growth form, forming thickets, mats, or clumps, the underground portions including a massive root crown, horizontal and vertical rhizomes, and an extensive root system; bark: thin, light brown and tinged with red, smooth or shallowly fissured. Leaves are deciduous, simple, alternate, ovate to nearly round, 2.5-3 cm long, with lateral, parallel veins in 8-13 pairs, the margins coarsely serrate or dentate to below middle or sometimes entire or with only a few small teeth at the top. Flowers are in short, dense, 5-15-flowered, upright racemes, the petals white, 1-2 cm long and strap-like, sepals more or less long-hairy on the inside, reflexing in age, stamens about 20, styles 5, ovary persistently tomentose at the top. Fruit are 6-11 mm long, smooth, purple-black, slightly gray-blue waxy, the pulp fleshy and sweet; seeds 4-10. The common name refers to the city in Saskatchewan, Canada, in the heart of the species' range.

Variation within the species: Considerable variation is recognized. Var. *alnifolia* occurs over the whole range of the species, except for California, where only var. *semiintegrifolia* is found. Other varieties include the following: var. *cusickii* (Fern.) C.L. Hitchc. and var. *humptulipensis* (G.N. Jones) C.L. Hitchc.

Distribution

Saskatoon is distributed along the Pacific coast from Alaska to California, eastward to Utah, Colorado, Iowa, and Minnesota, and Ontario and Quebec, north through the plains and prairies into the Northwest Territories (Keewatin and Mackenzie) of Canada. For current distribution, please consult the Plant Profile page for this species on the PLANTS Web site.

Adaptation

Saskatoon is common in lower-elevation coniferous forests but grows sporadically up to timberline. It also occurs in montane chaparral, mountain shrub, and the upper limits of pinyon-juniper communities. In grasslands, it mostly occurs in wooded draws, woodland interfaces, and riparian zones. It occurs in open to lightly shaded disturbed sites such as thickets, fencerows, clearings, and edges of woods, and it is conspicuous after disturbances such as fire, logging, or insect outbreak. Found at elevations of 50-3000 meters; flowering April-June; and fruiting (June-)July-August.

Establishment

Saskatoon reproduces from seed but more commonly by sprouting from the root crown and rhizomes and by layering. Seedlings can take up to 5 years before they start to produce fruit, but other starts may begin at 2-4 years old and with proper management can yield 8-10 tons of fruit per hectare. Flowers are produced almost every year, but good seed crops may be produced only every 3-5 years because of drought, spring frost, and/or juniper rust. Seeds require cold stratification to break dormancy and may remain viable for 10 or more years. Natural dispersal is by birds and mammals. One individual of saskatoon is known to have reached 85 years but the average age apparently may be closer to 20 years.

Management

Saskatoon in forests is fire-dependent, occurring in forests with fire regimes varying from frequent, low-severity fire (low-elevation forests) to infrequent, severe fire. It may persist in the understory for decades but eventually dies out with fire exclusion and canopy closure. After top-kill by light- to moderate-severity fire, saskatoon sprouts usually arise from the root crown or from shallowly buried rhizomes; sprouts arise from deeply buried rhizomes after even the most intense fire. Saskatoon cover and biomass production in western Montana may generally increase after fires in Douglas-fir/ninebark habitats, but browsing pressure from big game may slow the recovery. Seed production may resume soon after fire.

Cultivars, Improved and Selected Materials (and area of origin)

Saskatoon has been artificially crossed with other species of serviceberry, and similar hybridization and intergradation also occurs in the field, particularly between saskatoon and low serviceberry (*Amelanchier humilis* Wieg.). Cultivars sold as *Amelanchier alnifolia* may actually be hybrids between *A. alnifolia* and other species of *Amelanchier*. *Amelasorbus jackii* is a hybrid of saskatoon and Cascade Mountain-ash (*Sorbus scopulina*).

References

- Bishop, B.H. & S.H. Nelson 1980. *Propagation and transplanting of saskatoon (Amelanchier alnifolia Nutt.) softwood cuttings*. Canad. J. Pl. Sci. 60:883-890.
- Davidson, J.G.N. & R.G. St.-Pierre 1994. *Saskatoons*. HortScience 29(9):959-960.
- Howard, J.L. 1997. *Amelanchier alnifolia*. IN: D.G. Simmerman (compiler). *The fire effects information system* [Database]. USDA, Forest Service, Rocky Mountain Research Station, Intermountain Fire Sciences Laboratory, Missoula, Montana. <<http://www.fs.fed.us/database/feis/>>
- Jones, G.N. 1946. *American species of Amelanchier*. Illinois Biol. Monogr. 20(2):1-128.
- Laughlin, K.M., R.C. Smith, & .G. Askew 1996. *Juneberry for commercial and home use on the northern Great Plains*. Publ. H-938, North Dakota State University Extension Service, Fargo, North Dakota. <<http://www.ext.nodak.edu/extpubs/plantsci/hortcrop/h938w.htm>>
- Mazza, G. & .G. Davidson 1993. *Saskatoon berry: A fruit crop for the prairies*. Pp. 516-519, IN: J. Janick and J.E. Simon (eds.), *New crops*. Wiley, New York, New York. <<http://www.hort.purdue.edu/newcrop/proceedings1993/V2-516.html>>
- St-Pierre, R.G. 1992. *The development of native fruit species as horticultural crops in Saskatchewan*. HortScience 27(8):866, 947.
- St.-Pierre, R.G. 1997. *Growing Saskatoons - A manual for orchardists*. Completely rewritten Fifth Edition. Dept. of Horticulture Science, University of Saskatchewan, Saskatoon, Saskatchewan, Canada.

St.-Pierre, R.G. 1997. *Saskatoon*. Pp. 666-668 IN: *The Brooks and Olmo register of fruit & nut varieties*. Third Edition. ASHS Press, Alexandria, Virginia.

The Native Fruit Development Program 2000. *Registry of the genus Amelanchier: History & botany, cultivar overview, publications, plant & seed suppliers*. Dept. of Plant Sciences, University of Saskatchewan, Canada.
<<http://www.ag.usask.ca/departments/plsc/nfdp/amelanchier/index.html>>

Weir, B.J., R.N. Chibbar, & .G. St.-Pierre 1994. *RAPD fingerprinting cultivars of a native fruit species (Amelanchier alnifolia Nutt., Rosaceae), a new Canadian horticultural crop*. IN: P. Adams, J.S. Miller, E.M. Golenberg and J.E. Adams (eds.). *Conservation of plant genes II: Utilization of ancient and modern DNA*. Monogr. Syst. Bot. 48:141-148.

Weir, B.J., R.G. St.-Pierre, & R.N. Chibbar. 1997. *RAPD marker polymorphism among saskatoon cultivars, clones, and seedlings*. HortScience 32(6):1109-1113.

Prepared By

Guy Nesom
Formerly BONAP, North Carolina Botanical Garden,
University of North Carolina, Chapel Hill, North
Carolina

Species Coordinator

Lincoln Moore
USDA, NRCS, National Plant Data Center, Baton
Rouge, Louisiana

Edited: 17jan01 jsp;07feb03ahv; 30may06jsp

For more information about this and other plants, please contact your local NRCS field office or Conservation District, and visit the PLANTS Web site <<http://plants.usda.gov>> or the Plant Materials Program Web site <<http://Plant-Materials.nrcs.usda.gov>>

The U.S. Department of Agriculture (USDA) prohibits discrimination in all its programs and activities on the basis of race, color, national origin, sex, religion, age, disability, political beliefs, sexual orientation, and marital or family status. (Not all prohibited bases apply to all programs.) Persons with disabilities who require alternative means for communication of program information (Braille, large print, audiotape, etc.) should contact USDA's TARGET Center at 202-720-2600 (voice and TDD).

To file a complaint of discrimination write USDA, Director, Office of Civil Rights, Room 326-W, Whitten Building, 14th and Independence Avenue, SW, Washington, DC 20250-9410 or call

202-720-5964 (voice or TDD). USDA is an equal opportunity provider and employer.

[Read about Civil Rights at the Natural Resources Conservation Service.](#)